

**QUESTION 3: SOLUTION**

1. Using Coulomb's Law, we write the electric field at a distance  $r$  is given by

$$E_p = \frac{q}{4\pi\epsilon_0(r-a)^2} - \frac{q}{4\pi\epsilon_0(r+a)^2}$$

$$E_p = \frac{q}{4\pi\epsilon_0 r^2} \left( \frac{1}{\left(1-\frac{a}{r}\right)^2} - \frac{1}{\left(1+\frac{a}{r}\right)^2} \right) \dots\dots\dots(1)$$

Using binomial expansion for small  $a$ ,

$$E_p = \frac{q}{4\pi\epsilon_0 r^2} \left( 1 + \frac{2a}{r} - 1 + \frac{2a}{r} \right)$$

$$= + \frac{4qa}{4\pi\epsilon_0 r^3} = + \frac{qa}{\pi\epsilon_0 r^3} \dots\dots\dots(2)$$

$$= \frac{2p}{4\pi\epsilon_0 r^3}$$

2. The electric field seen by the atom from the ion is

$$\vec{E}_{ion} = -\frac{Q}{4\pi\epsilon_0 r^2} \hat{r} \dots\dots\dots (3)$$

The induced dipole moment is then simply

$$\vec{p} = \alpha \vec{E}_{ion} = -\frac{\alpha Q}{4\pi\epsilon_0 r^2} \hat{r} \dots\dots\dots (4)$$

From eq. (2)

$$\vec{E}_p = \frac{2p}{4\pi\epsilon_0 r^3} \hat{r}$$

The electric field intensity  $\vec{E}_p$  at the position of an ion at that instant is, using eq. (4),

$$\vec{E}_p = \frac{1}{4\pi\epsilon_0 r^3} \left[ -\frac{2\alpha Q}{4\pi\epsilon_0 r^2} \hat{r} \right] = -\frac{\alpha Q}{8\pi^2 \epsilon_0^2 r^5} \hat{r}$$

The force acting on the ion is

$$\vec{f} = Q\vec{E}_p = -\frac{\alpha Q^2}{8\pi^2 \epsilon_0^2 r^5} \hat{r} \dots\dots\dots (5)$$

The “-” sign implies that this force is attractive and  $Q^2$  implies that the force is attractive regardless of the sign of  $Q$ .

3. The potential energy of the ion-atom is given by  $U = \int_r^\infty \vec{f} \cdot d\vec{r}$  .....(6)

Using this,  $U = \int_r^\infty \vec{f} \cdot d\vec{r} = -\frac{\alpha Q^2}{32\pi^2 \epsilon_0^2 r^4}$  .....(7)

[Remark: Students might use the term  $-\vec{p} \cdot \vec{E}$  which changes only the factor in front.]

4. At the position  $r_{\min}$  we have, according to the Principle of Conservation of Angular Momentum,

$$mv_{\max} r_{\min} = mv_0 b$$

$$v_{\max} = v_0 \frac{b}{r_{\min}} \quad \dots\dots\dots (8)$$

And according to the Principle of Conservation of Energy:

$$\frac{1}{2}mv_{\max}^2 + \frac{-\alpha Q^2}{32\pi^2 \epsilon_0^2 r^4} = \frac{1}{2}mv_0^2 \quad \dots\dots\dots (9)$$

Eqs.(12) & (13):

$$\left(\frac{b}{r_{\min}}\right)^2 - \frac{\alpha Q^2 / \frac{1}{2}mv_0^2}{32\pi^2 \epsilon_0^2 b^4} \left(\frac{b}{r_{\min}}\right)^4 = 1$$

$$\left(\frac{r_{\min}}{b}\right)^4 - \left(\frac{r_{\min}}{b}\right)^2 + \frac{\alpha Q^2}{16\pi^2 \epsilon_0^2 mv_0^2 b^4} = 0 \quad \dots\dots\dots (10)$$

The roots of eq. (14) are:

$$r_{\min} = \frac{b}{\sqrt{2}} \left[ 1 \pm \sqrt{1 - \frac{\alpha Q^2}{4\pi^2 \epsilon_0^2 mv_0^2 b^4}} \right]^{\frac{1}{2}} \quad \dots\dots\dots (11)$$

[Note that the equation (14) implies that  $r_{\min}$  cannot be zero, unless  $b$  is itself zero.]

Since the expression has to be valid at  $Q = 0$ , which gives

$$r_{\min} = \frac{b}{\sqrt{2}} [1 \pm 1]^{\frac{1}{2}}$$

We have to choose “+” sign to make  $r_{\min} = b$

Hence,

$$r_{\min} = \frac{b}{\sqrt{2}} \left[ 1 + \sqrt{1 - \frac{\alpha Q^2}{4\pi^2 \epsilon_0^2 mv_0^2 b^4}} \right]^{\frac{1}{2}} \quad \dots\dots\dots(12)$$

5. A spiral trajectory occurs when (16) is imaginary (because there is no minimum distance of approach).

$r_{\min}$  is real under the condition:

$$1 \geq \frac{\alpha Q^2}{4\pi^2 \epsilon_0^2 m v_0^2 b^4}$$

$$b \geq b_0 = \left( \frac{\alpha Q^2}{4\pi^2 \epsilon_0^2 m v_0^2} \right)^{\frac{1}{4}} \dots\dots\dots (13)$$

For  $b < b_0 = \left( \frac{\alpha Q^2}{4\pi^2 \epsilon_0^2 m v_0^2} \right)^{\frac{1}{4}}$  the ion will collide with the atom.

Hence the atom, as seen by the ion, has a cross-sectional area  $A$ ,

$$A = \pi b_0^2 = \pi \left( \frac{\alpha Q^2}{4\pi^2 \epsilon_0^2 m v_0^2} \right)^{\frac{1}{2}} \dots\dots\dots (14)$$